

Numerical studies of the two- and three-dimensional gauge glass at low temperature

Helmut G. Katzgraber

Department of Physics, University of California, Davis, California 95616

(Dated: February 2, 2008)

We report results from Monte Carlo simulations of the two- and three-dimensional gauge glass at low temperature using parallel tempering Monte Carlo. In two dimensions, we find strong evidence for a zero-temperature transition. By means of finite-size scaling, we determine the stiffness exponent $\theta = -0.39 \pm 0.03$. In three dimensions, where a finite-temperature transition is well established, we find $\theta = 0.27 \pm 0.01$, compatible with recent results from domain-wall renormalization group studies.

PACS numbers: 75.50.Lk, 75.40.Mg, 05.50.+q

I. INTRODUCTION

The gauge glass is a model often used to describe the vortex glass transition in high-temperature superconductors. Still, there are areas which need to be understood. In two dimensions, there is an ongoing controversy as to whether a spin-glass transition occurs at finite temperature or not. In three dimensions, a finite-temperature transition is well established¹, although there is no consensus on the value of the stiffness exponent.

Recently, claims of evidence for a finite-temperature transition in two dimensions with $T_c = 0.22$ by Kim² using resistively shunted junction (RSJ) dynamics, and by Choi and Park³ who study the scaling of the spin-glass susceptibility via Monte Carlo simulations, have been made. In contrast, Granato⁴ and Hyman *et al.*⁵, who also use RSJ dynamics find evidence of a zero- T transition. In addition, Monte Carlo simulations by Fisher *et al.*⁶ and Reger and Young⁷ show evidence of a $T = 0$ transition, although the simulations were not performed at low enough temperatures.

Here we perform Monte Carlo simulations of the two- and three-dimensional gauge glass, using parallel tempering^{8,9}, to go to significantly lower temperatures than was possible in earlier work. In particular, we are now able to cover the temperature range in two dimensions where the claimed spin-glass transition^{2,3} takes place. We find strong evidence that $T_c = 0$.

In three dimensions, we study the gauge glass at very low but finite temperatures to provide a good estimate of the stiffness exponent θ . Earlier estimates, which were obtained from ground-state methods, are inconsistent. One group^{10,11,12,13} finds values consistent with $\theta \approx 0$, whereas another^{1,14,15,16,17} finds θ in the range $0.26 - 0.31$. We find that $\theta = 0.27 \pm 0.01$, which agrees with the results of Refs. 1, and 14,15,16,17.

II. MODEL, OBSERVABLES, AND METHOD

The Hamiltonian of the gauge glass is given by

$$\mathcal{H} = - \sum_{\langle i,j \rangle} \cos(\phi_i - \phi_j - A_{ij}), \quad (1)$$

where the sum ranges over nearest neighbors on a hypercubic lattice in D dimensions of size $N = L^D$, and ϕ_i represent the angles of the XY spins. Periodic boundary conditions are applied. The A_{ij} are quenched random variables uniformly distributed between $[0, 2\pi]$ ¹⁸. Because A_{ij} represent the line integral of the vector potential between sites i and j , we have the constraint that $A_{ij} = -A_{ji}$.

Traditionally, one uses the Binder ratio¹⁹ to estimate the critical temperature T_c . As for the gauge glass the Binder ratio cannot exceed unity¹, the splaying of the curves is small, and T_c is difficult to establish. In order to avoid this problem, we use a method introduced by Reger and Young¹⁰ in which one calculates the current I defined as the derivative of the free energy with respect to an infinitesimal twist to the boundaries, i.e.,

$$I(L) = \frac{1}{L} \sum_i \sin(\phi_i - \phi_{i+\hat{x}} - A_{i,i+\hat{x}}). \quad (2)$$

In this case, the twist is applied along the $\hat{x}$ -direction. As $[I(L)]_{\text{av}} = 0$, we actually calculate the root-mean-square current I_{rms} . Here $\langle \cdots \rangle$ represents a thermal average, whereas $[\cdots]_{\text{av}}$ represents a disorder average. Note that the current scales²⁰ as

$$I_{\text{rms}} = \tilde{I}(L^{1/\nu}(T - T_c)), \quad (3)$$

for $T_c > 0$, whereas for $T_c = 0$,

$$I_{\text{rms}} = L^{-1/\nu} \tilde{I}(L^{1/\nu} T). \quad (4)$$

Equation (4) indicates that the $T = 0$ stiffness exponent θ is negative and equal to $-1/\nu$ (as $I_{\text{rms}} \sim L^\theta$). Equation (3) shows that if $T_c > 0$, the curves for I_{rms} for different sizes *intersect* at the critical point, whereas Eq. (4) shows that if $T_c = 0$ the data *decrease* with increasing size at $T = 0$. For a finite-temperature transition we expect $\theta > 0$, since then the ordered state at $T = 0$ is “stiff” on large scales, and so will presumably resist small thermal fluctuations. On the other hand, for a zero-temperature transition, the system will then easily break up under the influence of thermal fluctuations and therefore $\theta < 0$.

We also have calculated the spin-glass susceptibility, χ_{SG} , defined by

$$\chi_{\text{SG}} = N[\langle q^2 \rangle]_{\text{av}}, \quad (5)$$

where q is the spin-glass order parameter:

$$q = \frac{1}{N} \sum_i^N \exp[i(\phi_i^\alpha - \phi_i^\beta)] . \quad (6)$$

Here, α, β are two replicas of the system with the same disorder. According to standard finite-size scaling the spin-glass susceptibility, defined in Eq. (5), behaves as

$$\chi_{\text{SG}} = L^{2-\eta} \tilde{\chi}_{\text{SG}}(L^{1/\nu}(T - T_c)) , \quad (7)$$

meaning that at criticality ($T = T_c$) it diverges with a power law, i.e., $\chi_{\text{SG}} \sim L^{2-\eta}$. Note that the power-law prefactor in Eq. (7) with an unknown exponent complicates the analysis of χ_{SG} compared with that for I_{rms} .

For the simulations, we use parallel tempering Monte Carlo^{8,9} as it allows us to study systems at lower temperatures than with conventional methods. For details about the implementation, equilibration tests, and detailed parameters of the simulations we refer the reader to Ref. 20. In two dimensions, the highest temperature is 1.058, whereas the lowest temperature is 0.13 for $L = 4$, 6, 8, 12, and 16 (for $L = 24$ the lowest temperature is 0.20). The number of temperatures $N_T = 30$ is chosen to give satisfactory acceptance ratios for the Monte Carlo moves between the temperatures ($N_T = 24$ for $L = 24$). In 3D the lowest temperature studied is 0.05 (note that $T_c \approx 0.45$)¹ whereas the highest temperature is 0.947, and $N_T = 53$ for $L = 3, 4, 5, 6$, and 8.

III. RESULTS FOR $D = 2$

Figure 1 shows a finite-size scaling plot of the data for I_{rms} in two dimensions according to Eq. (4). The plot shows that the data collapse if $L^{1/\nu}T$ is small for all values of L , and over the whole range of $L^{1/\nu}T$ for the largest sizes. The inset shows results of the unscaled data. At all temperatures, the data decrease with increasing L indicating, from Eq. (3), that T_c must be less than the range of temperatures studied. The data in Fig. 1 are therefore consistent with a zero-temperature transition, but with significant corrections to scaling at intermediate temperatures. We estimate the stiffness exponent to be $\theta = -0.39 \pm 0.03$. The above error bar is estimated by varying θ slightly until the data do not collapse well. This result is consistent with recent work of Akino and Kosterlitz¹⁴ who find $\theta = -0.36 \pm 0.01$. Choi and Park's³ parameters ($T_c = 0.22$, $1/\nu = 0.88$) yield *very* poor scaling, especially near the proposed T_c . In fact, we are unable to get a reasonable fit to the data for I_{rms} according to Eq. (3) for *any* finite T_c .

Figure 2 shows a scaling plot of the data for χ_{SG} according to Eq. (7) for $T_c = 0$ and $1/\nu = 0.39$, the same parameters found in the scaling of I_{rms} , together with $\eta = 0.0 \pm 0.1$, which is expected at a zero-temperature transition. The data at low T and for the largest sizes scale well, but the data away from this range show deviations. Allowing $1/\nu$ to vary, we find $1/\nu = 0.50 \pm 0.03$.




FIG. 1: Scaling of the root-mean-square current I_{rms} in two dimensions according to the form expected if $T_c = 0$, Eq. (4). We see acceptable scaling of the data at low temperatures. Deviations at higher T are presumably due to corrections to scaling. This plot is for $\theta \equiv -1/\nu = -0.39$. The inset shows I_{rms} as a function of T for different system sizes. At all temperatures, the data decrease with increasing L indicating, from Eq. (3), that if T_c is finite it must be less than 0.13.




FIG. 2: Scaling of χ_{SG} according to Eq. (7) with $T_c = 0$. The data for large sizes and low temperatures collapse with $\eta = 0$ and $1/\nu = 0.39$. The inset shows a scaling plot for the optimal value $1/\nu = 0.50$ (and $\eta = 0$). For these values of exponents, the collapse extends to a larger range of sizes.

The inset shows data for this optimal value, where only the $L = 4$ and 6 data are not part of the scaling function for all $L^{1/\nu}T$. The fact that the best values of $1/\nu$ are not precisely the same when obtained from χ_{SG} and I_{rms} presumably indicates that scaling is only valid for fairly low temperatures and large sizes, and that, despite our working at quite low temperatures, we have only a limited range of data which are fully in the scaling regime.

We also scale the data for χ_{SG} with Choi and Park's




FIG. 3: Effective stiffness exponent $\theta_{\text{eff}}(T)$ as a function of temperature in three dimensions for low temperatures. The data extrapolate linearly to $T = 0$.

parameters³. The data collapse is poor near T_c . The scaling of I_{rms} is also much worse, indicating that I_{rms} distinguishes between a finite T_c and $T_c = 0$ much better than χ_{SG} because of its simpler finite-size scaling form. An attempt to scale the data for χ_{SG} with $T_c = 0.13$, the lowest temperature simulated, shows that the best fit is obtained with $1/\nu = 0.68$ and $\eta = 0.19$. While the data scale acceptably well, the data for I_{rms} scale poorly.

According to Eq. (7) for $T = T_c$, the data for χ_{SG} should lie on a straight line at T_c . However, the data in the vicinity of $T = 0.22$, the transition temperature claimed by Kim² and Choi and Park,³ show a strong downward curvature²⁰ in our analysis, indicating that this is actually *above* T_c . Only around the lowest temperature where we have data, $T = 0.13$, is the curvature small, although it still greatly exceeds the error bars.

This indicates that $T_c < 0.13$, which is compatible with our data for I_{rms} .

IV. RESULTS FOR $D = 3$

Olson and Young¹ have investigated the critical region of the three-dimensional gauge glass obtaining a lower bound for the stiffness exponent of $\theta \geq 0.18$. Some previous results^{10,11,12,13} find θ in the range $0 \leq \theta \leq 0.077$ whereas others^{14,15,16,17} find a much larger value, $0.26 \leq \theta \leq 0.31$.

We can estimate θ from our data for I_{rms} since $I_{\text{rms}} \sim L^\theta$ when $L^{1/\nu}(T - T_c)$, the argument of the scaling function in Eq. (3), tends toward infinity. To obtain an estimate of θ , we perform a linear least-squares fit of $\ln(I_{\text{rms}})$ against $\ln(L)$ for each temperature in order to obtain an *effective* stiffness exponent $\theta_{\text{eff}}(T)$ which depends on the temperature. Figure 3 shows that $\theta_{\text{eff}}(T)$ can be fitted well to a linear form at low temperatures. Extrapolating to $T = 0$, we obtain $\theta = 0.27 \pm 0.01$, which is clearly positive and consistent with the results of Refs. 1,14,15,16,17.

V. CONCLUSIONS

Our results from Monte Carlo simulations of the two-dimensional gauge glass are consistent with a $T = 0$ transition with a stiffness exponent $\theta = -0.39 \pm 0.03$. These results are incompatible with the prediction made by Kim² and Choi and Park³ that $T_c = 0.22$. In three dimensions we report the first reliable estimate of the stiffness exponent from finite-temperature Monte Carlo simulations. We find $\theta = 0.27 \pm 0.01$, which agrees with the results of Refs. 1,14,15,16,17.

HGK acknowledges support from the National Science Foundation under Grant No. DMR 9985978 and would like to thank A. P. Young for a fruitful collaboration.

- ¹ T. Olson and A. P. Young, Phys. Rev. B **61**, 12467 (2000), (cond-mat/991229).
- ² B. J. Kim, Phys. Rev. B **62**, 644 (2000).
- ³ M. Y. Choi and S. Y. Park, Phys. Rev. B **60**, 4070 (1999).
- ⁴ E. Granato, Phys. Rev. B **58**, 11161 (1998).
- ⁵ R. A. Hyman, M. Wallin, M. P. A. Fisher, S. M. Girvin, and A. P. Young, Phys. Rev. B **51**, 15304 (1995).
- ⁶ M. P. A. Fisher, T. A. Tokuyasu, and A. P. Young, Phys. Rev. Lett. **66**, 2931 (1991).
- ⁷ J. D. Reger and A. P. Young, J. Phys. A **26**, 1067 (1993).
- ⁸ K. Hukushima and K. Nemoto, J. Phys. Soc. Jpn. **65**, 1604 (1996).
- ⁹ E. Marinari, in *Advances in Computer Simulation*, edited by J. Kertész and I. Kondor (Springer Verlag, Berlin, 1998), p. 50, (cond-mat/9612010).
- ¹⁰ J. D. Reger, T. A. Tokuyasu, A. P. Young, and M. P. A. Fisher, Phys. Rev. B **44**, 7147 (1991).
- ¹¹ M. J. P. Gingras, Phys. Rev. B **45**, 7547 (1992).

- ¹² J. M. Kosterlitz and M. V. Simkin, Phys. Rev. Lett. **79**, 1098 (1997).
- ¹³ J. Maucourt and D. R. Grempel, Phys. Rev. B **56**, 2572 (1997).
- ¹⁴ N. Akino and J. M. Kosterlitz (2002), (cond-mat/0203299).
- ¹⁵ M. Cieplak, J. R. Banavar, M. S. Li, and A. Khurana, Phys. Rev. B **45**, 786 (1992).
- ¹⁶ M. A. Moore and S. Murphy, Phys. Rev. B **50**, 3450 (1994).
- ¹⁷ J. M. Kosterlitz and N. Akino, Phys. Rev. Lett. **81**, 4672 (1998).
- ¹⁸ H. G. Katzgraber and A. P. Young, Phys. Rev. B **64**, 104426 (2001), (cond-mat/0105077).
- ¹⁹ K. Binder, Phys. Rev. Lett. **47**, 693 (1981).
- ²⁰ H. G. Katzgraber and A. P. Young, Phys. Rev. B **66**, 224507 (2002), (cond-mat/0205206).